

A.

Ref BASIS: 17958

N° d'accès Derwent: 91-356402 [49]

Titre Derwent: Cell stimulants for external use e.g. alopecia, acne etc. contains dextran, glucose, mutan, lentinan, sodium and potassium chloride and calcium hydrochloride

TEXTO_Mots-Clés Fapp: 14 DOMAINE ALOPECIE ACNE DERMATITE SOIN FORMULATION DEXTRANE GLUCOSE MUTAN LENTINAN NaCl KCl CALCIUM

Domaine Principal: Antichute et repousse (cheveux)

Déposants: JAPAN FINE CHEM CO LTD; JAPAN FINE CHEM KK

Inventeurs: YAMADA A; YAMADA H

Priorité (N° + date): 90JP-034720 900215

Dépôt (N° + date): 90JP-034720 900215; 91EP-401785 910628; 91AU-078499 910618; 91CA-044795 910617; 91AU-078499 910618 [Previous Publ. AU9178499]

Publication (N° + date): J03240730 A 911028 DW9149 000pp; US5126135 A 920630 DW9229 A61K-007/48 004pp; EP-520112 A1 921230 DW9301 A61K-007/48 Eng 007pp!; AU9178499 A 930107 DW9308 A61K-007/48 000pp!; CA2044795 A 921218 DW9310 A61K-007/48 000pp!; AU-639793 B 930805 DW9338 A61K-007/48 000pp!

Résumé (+ Use/Advantage): J03240730 Cell stimulants contain 5-30 wt.% of dextran, 5-30 wt.% of glucose, 5-30% of mutan, 5-30 wt.% of lentinan, 0.1-1 wt.% of sodium chloride, 0.1-1 wt.% of calcium hydrochloride and 0.1-1 wt.% of potassium chloride. USE | The drug is used for alopecia, acne, dermatitis, tinea pedis, stomatitis and pollinosis. The drug stimulates the cell activity due to osmotic effect. (4pp Dwg.No.0/0); EP-520112 Cell stimulants contain 5-30 wt.% of dextran, 5-30 wt.% of glucose, 5-30% of mutan, 5-30 wt.% of lenthynan, 0.1-1 wt.% of sodium chloride, 0.1-1 wt.% of calcium chloride and 0.1-1 wt.% of potassium chloride. USE | The drug is used for alopecia, acne, dermatitis, tinea pedis, stomatitis and pollinosis. The drug stimulates the cell activity due to osmotic effect. (Dwg.0/0); US5126135 Compsn. based on an aq. phase comprises dextran glucose, mutan, lentinan, NaCl, KCl and CaCl2 dissolved in the aq. phase. The concn. of dextran, glucose, lentinan are each 50-30% of the wt. of the aq. phase and the concn. of NaCl, KCl and CaCl2 are each between 0.1-1% of the wt. of aq. phase. The aq. phase does not contain any oily material and is at 50-100 deg.C for the dissolution. USE | Compsn. is a dermatological compsn. of cosmetics or medicaments for external application to skin and to oral or mucous membrane. Used to regenerate and revitalise damaged cells by equilibrating electrochemical ion gradients across cell membrane and by osmotic pressure on inner face of cell wall. (Dwg.0/0)

Use / Advantage: J03240730 < USE > The drug is used for alopecia, acne, dermatitis, tinea pedis, stomatitis and pollinosis. The drug stimulates the cell activity due to osmotic effect. (4pp Dwg.No.0/0); EP-520112 < USE > The drug is used for alopecia, acne, dermatitis, tinea pedis, stomatitis and pollinosis. The drug stimulates the cell activity due to osmotic effect. (Dwg.0/0);

20 May 1999